

Supplemental Table 2. Secreted proteins increased in both cortical collecting ducts (CCDs) and cortical thick ascending limbs of Henle's loop (CTALs) three hours after unilateral ureteral obstruction (UUO).

Gene symbol	Annotation	CCD Log₂ (UUO/ Sham)	CTAL Log₂ (UUO/ Sham)
<i>Afm</i>	afamin	4.073	1.752
<i>Angptl4</i>	angiopoietin-related protein 4	7.400	8.014
<i>Apoc2</i>	apolipoprotein C-II	2.529	2.330
<i>ApoE</i>	apolipoprotein E	1.663	1.046
<i>Apom</i>	apolipoprotein M	2.902	3.823
<i>Cela1</i>	chymotrypsin-like elastase family member 1	1.521	1.235
<i>Hbegf</i>	proheparin-binding EGF-like growth factor	2.998	3.706
<i>Lcn2</i>	neutrophil gelatinase-associated lipocalin	3.766	2.351
<i>Napsa</i>	napsin-A	2.378	2.410
<i>Pm20d1</i>	N-fatty-acyl-amino acid synthase/hydrolase PM20D1	2.214	2.622
<i>Serpinf1</i>	pigment epithelium-derived factor	2.343	1.750
<i>Serpinf2</i>	alpha-2-antiplasmin	2.938	1.716
<i>Tff3</i>	trefoil factor 3	2.361	2.385
<i>Timp1</i>	metalloproteinase inhibitor 1	2.489	1.040